

## Appendix B

### Summary of Human Clinical Trials of Foodborne Pathogens

Potential Surrogate Pathogens for <i>E. coli</i> O157:H7 (strain or serotype)	Number of Dose Groups	Total Number of Human Volunteers	Lowest Dose Tested and with Illness	Reference
1. <i>S. dysenteriae</i> (M131) <sup>a</sup>	4	30 <sup>b</sup>	10, 10	Levine et al. 1973
2. <i>S. dysenteriae</i> (A-1) <sup>c</sup>	2	10	2x10 <sup>2</sup> , 2x10 <sup>2</sup>	Levine et al. 1973
3. <i>S. flexneri</i> (2457T) <sup>d</sup>	5	43	10 <sup>4</sup> , 10 <sup>4</sup>	DuPont et al. 1969
4. <i>S. flexneri</i> (2457T)	4	197	10 <sup>5</sup> , 10 <sup>5</sup>	DuPont et al. 1972
5. <i>S. sonnei</i> (53G) <sup>e</sup>	1	20, 38	500, 500	DuPont et al. 1989
6. Enteropathogenic <i>E. coli</i> wild type (O111:NM, B1718) <sup>f</sup> + bicarbonate	3	13 <sup>g</sup>	5x10 <sup>8</sup> , 5x10 <sup>8</sup>	Bieber et al. 1998
7. Enteropathogenic <i>E. coli</i> (O142:H6) <sup>h</sup> + bicarbonate	3	15 <sup>i</sup>	10 <sup>6</sup> , 10 <sup>6</sup>	Levine et al. 1978
8. Enteropathogenic <i>E. coli</i> (O128:H6) <sup>j</sup> + bicarbonate	3	15	Avirulent at 10 <sup>6-10</sup>	Levine et al. 1978
9. Enteropathogenic <i>E. coli</i> (O127:H6) <sup>k</sup> + bicarbonate	2	9	10 <sup>6</sup> , 10 <sup>10</sup>	Levine et al. 1978
10. Enterotoxigenic <i>E. coli</i> (O78:H11) <sup>l</sup> + bicarbonate	2	14 <sup>m</sup>	10 <sup>6</sup> , 10 <sup>6</sup>	Evans et al. 1978
11. Enterotoxigenic <i>E. coli</i> (non-typable) <sup>n</sup>	3	14 <sup>o</sup>	10 <sup>6</sup> , 10 <sup>8</sup>	Levine et al. 1977
12. Enterotoxigenic <i>E. coli</i> (O148:H28) <sup>p</sup>	2	17 <sup>q</sup>	10 <sup>6</sup> , 10 <sup>6</sup>	Levine et al. 1979
13. Enterotoxigenic <i>E. coli</i> (O25:NM) <sup>r</sup>	1	6 <sup>o</sup>	10 <sup>9</sup> , 10 <sup>9</sup>	Levine et al. 1979

Potential Surrogate Pathogens for <i>E. coli</i> O157:H7 (strain or serotype)	Number of Dose Groups	Total Number of Human Volunteers	Lowest Dose Tested and with Illness	Reference
14. Infant diarrheal <i>E. coli</i> (O111, B4, H)	4	46 <sup>s</sup>	10 <sup>6</sup> , 10 <sup>6</sup>	June et al. 1953
15. Infant diarrheal <i>E. coli</i> (O111, B4, H)	4	46 <sup>s</sup>	10 <sup>8</sup> , 10 <sup>8</sup>	June et al. 1953
16. Infant diarrheal <i>E. coli</i> (O111, B5)	1	1 <sup>t</sup>	10 <sup>8</sup> , 10 <sup>8</sup>	Ferguson and June 1952, citing Neter 1950
17. Infant diarrheal <i>E. coli</i> (O111, B5)	1	6	10 <sup>9</sup> , 10 <sup>9</sup>	Ferguson and June 1952, citing Kirby 1950
18. Infant diarrheal <i>E. coli</i> (O111, B5)	1	3	Avirulent at 10 <sup>10</sup>	Ferguson and June 1952, citing Kirby 1950
19. Commensal <i>E. coli</i>	2	19	Avirulent at ~10 <sup>10</sup>	June et al. 1953
20. Commensal <i>E. coli</i> <sup>u</sup> + bicarbonate	1	4	Avirulent at 10 <sup>10</sup>	Levine et al. 1978

<sup>a</sup>Isolated from feces of patient in Guatemala with severe dysentery from 1970 pandemic and administered in milk.

<sup>b</sup>Fasting male prison volunteers.

<sup>c</sup>Isolated from feces of patient in Guatemala with mild dysentery and administered in milk.

<sup>d</sup>Isolated from feces of patient in Japan and administered in milk.

<sup>e</sup>Isolated from feces of 5-year-old patient in Japan and administered in milk.

<sup>f</sup>Isolated and administered in phosphate buffered saline with sodium bicarbonate.

<sup>g</sup>Fasting volunteers, 18 to 48 years of age.

<sup>h</sup>Isolate infant diarrheal strain from UK hospitals (Glasgow, E851/71) and administered with bicarbonate in saline; virulent at each of 3 doses administered.

<sup>i</sup>Healthy adult volunteers, mean age 24 years, 90-minute fast pre- and post-treatment.

<sup>j</sup>Isolate infant diarrheal strain from UK hospitals (Teesside, E74/68) and administered with bicarbonate in saline; avirulent in 15 healthy adults tested.

<sup>k</sup>Isolate infant diarrheal strain from UK residential nursery (Taunton, E2348/69) and administered with bicarbonate in saline; virulent at 1 of 2 doses administered.

<sup>l</sup>Isolated from severe non-Vibrio cholera case in Bangladesh.

<sup>m</sup>Male and female student volunteers, mean age 23.

<sup>n</sup>Isolated from physician traveling in Mexico with traveler's diarrhea and administered in milk with 2.5 hour fast intervals before and after treatment.

<sup>o</sup>Student volunteers, 18 to 29 years of age.

<sup>p</sup>Isolated from U.S. soldier in Vietnam with diarrhea and administered with bicarbonate in buffer.

<sup>q</sup>Healthy adult volunteers, mostly students, mean age 25, range 18 to 41 years of age.

<sup>r</sup>Isolated from physician traveling in Mexico with traveler's diarrhea and administered in milk with 2.5 hour fast intervals before and after treatment.

<sup>s</sup>Presumably fasting male prison volunteers, 16 to 48 years of age.

<sup>t</sup>Two-month old infant administered 10<sup>8</sup> organisms and developed diarrhea and weight loss within 24 hours.

<sup>u</sup>Nonpathogenic isolate from healthy laboratory scientist, administered in bicarbonate at 10<sup>10</sup> in saline; avirulent in all four volunteers.

## REFERENCES

- Bieber, D., S.W. Ramer, C.-Y. Wu, W.J. Murray, T. Tobe, R. Fernandez, and G.K. Schoolnik. 1998. Type IV pili, transient bacterial aggregates, and virulence of enteropathogenic *Escherichia coli*. *Science* 280:2114-2118.
- Dupont, H.L., R.B. Hornick, A.T. Dawkins, M.J. Snyder, and S.B. Formal. 1969. The response of man to virulent *Shigella flexneri* 2a. *J Infect Dis* 119:296-299.
- Dupont, H.L., R.B. Hornick, M.J. Snyder, J.P. Libonati, S.B. Formal, and E.J. Gangarosa. 1972. Immunity in Shigellosis II. Protection induced by oral live vaccine or primary infection. *J Infect Dis* 125(1):12-16.
- DuPont, H.L., M.M. Levine, R.B. Hornick, and S.B. Formal. 1989. Inoculum size in shigellosis and implications for expected mode of transmission. *J Infect Dis* 159(6):1126-1128.
- Evans, D.G., D.J. Evans, Jr., W.S. Tjoa, and H.L. Dupont. 1978. Detection and characterization of colonization factor of enterotoxigenic *Escherichia coli* isolated from adults with diarrhea. *Infect Immun* 19:727-736.
- Ferguson, W.W., and R.C. June. 1952. Experiments on feeding adult volunteers with *Escherichia coli* 111, B5, a coliform organism associated with infant diarrhea. *Am J Hyg* 55(2):155-169.
- June, R.C., W.W. Ferguson, and M.T. Worfel. 1953. Experiments in feeding adult volunteers with *Escherichia coli* 55, B5, a coliform organism associated with infant diarrhea. *Am J Hyg* 57:222-236.
- Levine, M.M., H.L. DuPont, S.B. Formal, R.B. Hornick, A. Takeuchi, E.J. Gangarosa, M.J. Snyder, and J.P. Libonati. 1973. Pathogenesis of *Shigella dysenteriae* 1 (Shiga) dysentery. *J Infect Dis* 127(3):261-270.
- Levine, M.M., E.S. Caplan, D. Waterman, R.A. Cash, R.B. Hornick, and M.J. Snyder. 1977. Diarrhea caused by *Escherichia coli* that produce only heat-stable enterotoxin. *Infect Immun* 17(1):78-82.
- Levine, M.M., E.J. Bergquist, D.R. Nalin, D.H. Waterman, R.B. Hornick, C.R. Young, and S. Sotman. 1978. *Escherichia coli* strains that cause diarrhoea but do not produce heat-labile or heat-stable enterotoxins and are non-invasive. *Lancet* 1(8074):1119-1122.
- Levine, M.M., D.R. Nalin, D.L. Hoover, E.J. Bergquist, R.B. Hornick, and C.R. Young. 1979. Immunity to enterotoxigenic *Escherichia coli*. *Infect Immun* 23(3):729-736.